

TEREBRIDAE
ex coll. Don Pisor

Author: Yves Terryn

Don Pisor coll.



Trawled, Taiwan. 98.0 mm

linked to
lima

2010 - singleton DP
2016 - TY, SH, MC



Trawled, Taiwan. 97.3 mm

linked to
fenestrata

2010 - singleton DP
2016 - TY, SH, MC

cumingii



B&C, 1987



lectotype



???

fenestrata



actual lectotype,
but as paralectotype in
NHM coll.



lectotype of *caelata*

fenestrata



actual paralectotype,
but as lectotype in
NHM coll.



actual lectotype,
but as paralectotype in
NHM coll.

anilis



actual paralectotype of
fenestrata,
but as lectotype in
NHM coll.



holotype of
rubrobrunnea

anilis



holotype of
anilis



holotype of
circinata

lima



holotype of *lima*



lectotype of *cumingii*

lima



holotype of *lima*



holotype of *loebbeckeana*



loebbeckeana



holotype of *loebbeckeana*



pretiosa



fujitai


loebbeckeana



holotype of *loebbeckeana*



commaculata



holotype of *myuros*

cossignanii



holotype of *loebbeckeana*



holotype of *myuros*



holotype of *cossignanii*

???



???



holotype of *straminea*



holotype of
anilis

fenestrata



actual lectotype,
but as paralectotype in
NHM coll.



probable syntype of *torquata*,
'similar' to original drawing

boucheti



holotype MNHN



paratype NHM



paratypes LACM



TYPES

Bratcher & Cernohorsky, 1987

anilis
straminea
circinata
rubrobrunnea
serotina
commaculata
myuros
cossignanii
lima
loebbeckeana
pretiosa
fujitai

cumingii
boucheti
fenestrata
 = *caelata*
 = *torquata*
punctatostriata
 = *pallida*

9 species, 10 taxa in synonymy

(2016)

anilis
straminea
circinata
rubrobrunnea
commaculata
myuros
cossignani
lima
loebbeckeana
pretiosa
fujitai
cumingii
boucheti
fenestrata
 = *caelata*
torquata
punctatostriata
 = *pallida*

serotina nomen nudum - type lost, description/figure too vague

15 (14) species, 1 (2) taxon in synonymy, 1 taxon excluded

fenestrata
= *caelata*



cumingii / *lima* / *loebbeckeana*



myuros / cossignanii
= ?



punctatostriata
= *pallida*



straminea



torquata



probably syntypes + specimens
yet prob. syntypes differ from original drawing



specimens closer to original
drawing and description


anilis



anilis - circinata (specimens < 30 mm)




others (specimens > 35 mm)



smooth form (B&C, 1987) = ??? 2



???
amanda group




juveniles of ??? 1

species 1 aff. lima



species 2 aff. torquata
see 8

Vanuatu



same trawl S Japan

species 3 aff. torquata
see 8




Balicasag, MNHN, + 250 m
see Xenophora pallidula

species 4 - amanda s. l. group

"anilis ??? form"

punctatostriata

"anilis smooth form"



species 5
Vietnam



Vietnam

species 10



India

Vietnam

Philippines

species 9 'old cumingii' = ??? 1 at the beginning



species 9 'old cumingii' = ??? 1 at the beginning



species 6 a



Madagascar +250 m

species 6 b



S Japan

species 6 b



India


Philippines

species 7 - donpisorii





???
boucheti
one of the LACM paratypes

species 8 aff. torquata



boucheti





boucheti

species 12-14

species 11 = 2 LACM paratypes



species 15



species 16





Xenophora pallidula

sps of cf. *torquata*



species 8 aff. *torquata*



species aff. *donpisoni*



neglecta



swobodai



cf. *subtextilis*



sp. nov.



sp. nov. in prep.

